CONTENTS

(Abstracts/contents lists published in: Chem. Abstr.; Curr. Contents; Phys. Chem. Earth Sci., Life Sci.)

Dedication: Professor Walter T. J. Morgan xi

Structure elucidation of blood group B-like and I-active ceramide eicosa- and pentacosasaccharides from rabbit erythrocyte membranes by combined gas chromatography–mass spectrometry; electron-impact and fast-atom-bombardment mass spectrometry; and two-dimensional correlated, relayed-coherence transfer, and nuclear Overhauser effect 500-MHz ^{1}H-n.m.r. spectroscopy
P. HANFLAND, M. KORDOWICZ, J. PETER KATALINIC, H. EGGE (Bonn, F.R.C.), J. DABROWSKI, AND U. DABROWSKI (Heidelberg, F.R.G.) 1

Biosynthesis of cancer-associated sialyl-X antigen by a (1→3)-α-L-fucosyltransferase of human amniotic fluid
F.-G. HANISCH, A. MITSAKOS, H. SCHROTEN, AND G. UHLENBRUCK (Cologne, F.R.G.) . 23

Structural studies on oncofetal carbohydrate antigens (CA 19-9, CA 50, and CA 125) carried by *O*-linked sialyloligosaccharides on human amniotic mucins
F.-G. HANISCH, G. UHLENBRUCK (Cologne, F.R.G.), J. PETER KATALNIC, AND H. EGGE (Bonn, F.R.G.) 29

Isolation and partial purification of lipid-linked oligosaccharides from calf pancreas
J. BADET AND R. W. JEANLOZ (Boston, MA, U.S.A.) 49

500-MHz ^{1}H-n.m.r. and conformational studies of fucosyl-oligosaccharides recognised by monoclonal antibodies with specificities related to Le^a, Le^b, and SSEA-1
E. F. HOUNSELL, N. J. JONES, H. C. GOOI, T. FEIZI, A. S. R. DONALD (Middlesex, Gt. Britain) and J. FEENEY (London, Gt. Britain) 67

Separation of human milk oligosaccharides by recycling chromatography. First isolation of lacto-*N-neo*-difucohexaose II and 3′-galactosyllactose from this source
A. S. R. DONALD (Middlesex, Gt. Britain) AND J. FEENEY (London, Gt. Britain) . . 79

Mucus glycoprotein gels. Role of glycoprotein polymeric structure and carbohydrate side-chains in gel-formation
L. A. SELLERS, A. ALLEN (Newcastle upon Tyne, Gt. Britain) E. R. MORRIS, AND S. B. ROSS-MURPHY (Bedford, Gt. Britain) 93

Structures of the eight- to nine-sugar glycolipids of human blood group A erythrocytes
M. E. BREIMER, H. KARLSSON, K.-A. KARLSSON, K. NILSON, B. E. SAMUELSSON, AND N. STRÖMBERG (Göteborg, Sweden) 111

^{1}H-N.m.r. analysis of Type-2 chain lacto-gangliosides. Conformation of structure of a novel cancer-associated fucoganglioside, α-NeuAc-(2→6)-β-D-Gal*p*-(1→4)-β-D-Glc*p*-NAc-(1→3)-β-D-Gal*p*-(1→4)-[α-L-Fuc*p*-(1→3)]-β-D-Glc*p*NAc-(1→3)-β-D-Gal*p*-(1→4)-β-D-Glc*p*-(1→1)-Cer ($VI^6NeuAcIII^3FucnLc_6Cer$)
S. B. LEVERY, E. NUDELMAN, R. KANNAGI, F. W. SYMINGTON, N. H. ANDERSEN, H. CLAUSEN, M. BALDWIN, AND S. HAKOMORI (Seattle, WA, U.S.A.) 121

Oligosaccharides from faeces of a blood-group B, breast-fed infant
H. SABHARWAL, B. NILSSON, M. A. CHESTER, F. LINDH, G. GRÖNBERG, S. SJÖBLAD, AND A. LUNDBLAD (Lund, Sweden) 145

CONTENTS

(Abstracts/contents lists published in: Chem. Abstr.; Curr. Contents; Phys. Chem. Earth Sci., Life Sci.)

Dedication: Professor Walter T. J. Morgan xi

Structure elucidation of blood group B-like and I-active ceramide eicosa- and pentacosasaccharides from rabbit erythrocyte membranes by combined gas chromatography–mass spectrometry; electron-impact and fast-atom-bombardment mass spectrometry; and two-dimensional correlated, relayed-coherence transfer, and nuclear Overhauser effect 500-MHz ^{1}H-n.m.r. spectroscopy
P. HANFLAND, M. KORDOWICZ, J. PETER KATALINIC, H. EGGE (Bonn, F.R.C.), J. DABROWSKI, AND U. DABROWSKI (Heidelberg, F.R.G.) 1

Biosynthesis of cancer-associated sialyl-X antigen by a (1→3)-α-L-fucosyltransferase of human amniotic fluid
F.-G. HANISCH, A. MITSAKOS, H. SCHROTEN, AND G. UHLENBRUCK (Cologne, F.R.G.) . 23

Structural studies on oncofetal carbohydrate antigens (CA 19-9, CA 50, and CA 125) carried by *O*-linked sialyloligosaccharides on human amniotic mucins
F.-G. HANISCH, G. UHLENBRUCK (Cologne, F.R.G.), J. PETER KATALNIC, AND H. EGGE (Bonn, F.R.G.) 29

Isolation and partial purification of lipid-linked oligosaccharides from calf pancreas
J. BADET AND R. W. JEANLOZ (Boston, MA, U.S.A.) 49

500-MHz ^{1}H-n.m.r. and conformational studies of fucosyl-oligosaccharides recognised by monoclonal antibodies with specificities related to Le^a, Le^b, and SSEA-1
E. F. HOUNSELL, N. J. JONES, H. C. GOOI, T. FEIZI, A. S. R. DONALD (Middlesex, Gt. Britain) and J. FEENEY (London, Gt. Britain) 67

Separation of human milk oligosaccharides by recycling chromatography. First isolation of lacto-*N-neo*-difucohexaose II and 3′-galactosyllactose from this source
A. S. R. DONALD (Middlesex, Gt. Britain) AND J. FEENEY (London, Gt. Britain) . . 79

Mucus glycoprotein gels. Role of glycoprotein polymeric structure and carbohydrate side-chains in gel-formation
L. A. SELLERS, A. ALLEN (Newcastle upon Tyne, Gt. Britain) E. R. MORRIS, AND S. B. ROSS-MURPHY (Bedford, Gt. Britain) 93

Structures of the eight- to nine-sugar glycolipids of human blood group A erythrocytes
M. E. BREIMER, H. KARLSSON, K.-A. KARLSSON, K. NILSON, B. E. SAMUELSSON, AND N. STRÖMBERG (Göteborg, Sweden) 111

^{1}H-N.m.r. analysis of Type-2 chain lacto-gangliosides. Conformation of structure of a novel cancer-associated fucoganglioside, α-NeuAc-(2→6)-β-D-Gal*p*-(1→4)-β-D-Glc*p*-NAc-(1→3)-β-D-Gal*p*-(1→4)-[α-L-Fuc*p*-(1→3)]-β-D-Glc*p*NAc-(1→3)-β-D-Gal*p*-(1→4)-β-D-Glc*p*-(1→1)-Cer ($VI^6NeuAcIII^3FucnLc_6Cer$)
S. B. LEVERY, E. NUDELMAN, R. KANNAGI, F. W. SYMINGTON, N. H. ANDERSEN, H. CLAUSEN, M. BALDWIN, AND S. HAKOMORI (Seattle, WA, U.S.A.) 121

Oligosaccharides from faeces of a blood-group B, breast-fed infant
H. SABHARWAL, B. NILSSON, M. A. CHESTER, F. LINDH, G. GRÖNBERG, S. SJÖBLAD, AND A. LUNDBLAD (Lund, Sweden) 145

A method for reductive cleavage of *N*-glycosylamide carbohydrate–peptide bond
L. M. Likhosherstov, O. S. Novikova, V. E. Piskarev, E. E. Trusikhina, V. A. Derevitskaya, and N. K. Kochetkov (Moscow, U.S.S.R.) 155

The carbohydrate chains of influenza virus hemagglutinin
N. P. Arbatsky, V. A. Derevitskaya, A. O. Zheltova, N. K. Kochetkov, L. M. Likhosherstov, S. N. Senchenkova, and D. V. Yurtov (Moscow, U.S.S.R.) . . 165

Studies on the chemical modification of soybean agglutinin
N. N. Desai, A. K. Allen, and A. Neuberger
(London, Gt. Britain) 183

Structure of the capsular K3 antigen of *Escherichia coli* O4:K3:H4, a polysaccharide containing a 4-deoxy-2-hexulosonic acid
T. Dengler, K. Himmelspach, B. Jann, and K. Jann (Freiburg, F.R.G.) 191

Effect of endoglycosidase F-peptidyl *N*-glycosidase F preparations on the surface components of the human erythrocyte
M. J. A. Tanner, D. J. Anstee, G. Mallinson, K. Ridgwell, P. G. Martin, N. D. Avent, and S. F. Parsons, (Bristol, Gt. Britain) 203

Carbohydrate-binding specificities of five lectins that bind to *O*-glycosyl-linked carbohydrate chains. Quantitative analysis by frontal-affinity chromatography
S. Sueyoshi, T. Tsuji, and T. Osawa (Tokyo, Japan) 213

Removal of ester-bound fatty acyl residues from *Salmonella* lipopolysaccharide by enzyme preparations from *Acanthamoeba castellanii*
W. Drozanski, A. E.-H. A. Moustafa, O. Lüderitz, and O. Westphal (Freiburg, F.R.G.) 225

The comparison of monoclonal antibodies specific for two general epitopes located on *Citrobacter* O36 lipopolysaccharide
M. Kulakowska, C. Lugowski, E. Romanowska (Wrocław, Poland) and D. R. Bundle (Ottawa, Ontario, Canada) 233

Immunochemical studies on the *N*-acetyllactosamine β-(1→6)linked trisaccharide specificity of *Ricinus communis* agglutinin
A. M. Wu, S. Sugii (College Station, TX, U.S.A.), F. G. Gruezo and E. A. Kabat (New York, NY, U.S.A.) 243

Substrate recognition by UDP-*N*-acetyl-α-D-galactosamine: polypeptide *N*-acetyl-α-D-galactosaminyltransferase. Effects of chain length and disulphide bonding of synthetic peptide substrates
R. C. Hughes, A. F. Bradbury, and D. G. Smyth (London, Gt. Britain) 259

Blood group Tn-active macromolecules from human carcinomas and erythrocytes. Characterization of and specific reactivity with mono- and poly-clonal anti-Tn antibodies induced by various immunogens
G. F. Springer, E. V. Chandrasekaran, P. R. Desai, and H. Tegtmeyer (Evanston, IL, U.S.A.) 271

The binding of the Lewis-a human blood group determinant by two hybridoma monoclonal anti-Le^a antibodies
R. U. Lemieux, O. Hindsgaul, P. Bird, S. Narasimhan (Edmonton, Alberta, Canada), and W. W. Young, Jr. (Charlottesville, VA, U.S.A.) 293

(Continued overleaf)

(contents continued)

The effect of glycosidases on the survival of rat erythrocytes in circulation
K. A. GUTOWSKI, D. A. LINSEMAN, AND D. AMINOFF (Ann Arbor, MI, U.S.A.) . . 307

Notes

Oxidation products of ovarian cyst mucins. Characterisation by density-gradient analysis
J. M. CREETH (Bristol, Gt. Britain) 314

Blood-group active proteoglycan containing 3-*O*-methylrhamnose (acofriose) from young plants of *Osmunda japonica*
T. AKIYAMA, K. TANAKA, S. YAMAMOTO, AND S. ISEKI (Tokyo, Japan) 320

Structural analysis of the preponderant high-mannose oligosaccharide of human Tamm-Horsfall glycoprotein
F. DALL'OLIO, F. J. J. DE KANTER, D. H. VAN DEN EIJNDEN, AND F. SERAFINI-CESSI (Amsterdam, The Netherlands) 327

Analysis of a mouse monoclonal antibody detecting the H Type 1 blood group determinant
S. WELT, B. W. T. YIN, K. FURUKAWA, K.-A. KARLSSON, L. J. OLD, AND K. O. LLOYD (Göteborg, Sweden) 333

Announcement c1

Author index c3

Subject index c5